FULVIC ACIDS AT 80% SOIL IMRPOVER

FulvicProtect is 100% of vegetable origin and is obtained through a specialized process within humic substances. It stimulates absorption and penetration through the active transport of nutrients at the membrane level of leaf and root cells. It has a high cation exchange capacity.

BENEFITS

- Improves soil structure.
- High cation exchange capacity.
- Increases nutrient absorption via foliar and root.
- Strengthens roots and improves plant vigor.

BIOLOGICAL COMPOSITION

Fulvic Acids	80%
Inert Material	20%

AGRONOMIC USE

FulvicProtect guarantees a 100% organic and high-quality product that can be used for the following crops, although this recommendation is not limited to other agricultural crops: **Vegetables in general:** Cabbage, cauliflower, broccoli, compass cabbage, radish. **Fruit trees in general:** Avocado, citrus, peach, guava, mango, coconut, banana, papaya. **Solanaceae**: Tomato, chili, potato, eggplant, pepper. **Cucurbits**: Cucumber, melon, watermelon, pumpkins. **Seeds:** corn and sorghum, wheat, barley, and oats. **Berries**: blueberry, raspberry, blackberry, blackberry. **Other crops:** Celery, asparagus, lettuce, onion, chives, carrot, walnut, pineapple, vine, sugar cane, cotton, agaves, oil palm.

APPLICATION METHOD

FulvicProtect is applied by injecting it into the root with a pump or directly into the irrigation system, as well as via foliar application. Always ensure that the equipment used has not previously contained a pesticide or herbicide. Clean the pump thoroughly before use, preferably using neutral powder soap. It is recommended to dilute 250 g of **FulvicProtect** in the cleanest possible natural water per hectare for crops weighing 1-9 kg. For crops weighing 10-50 kg, use 1 kg per hectare, and for 60-100 kg, use 1.5 kg per hectare of **FulvicProtect**. Do not dilute in more than 200 L of water for every 250 g of **FulvicProtect**, as minimal doses are needed to create a favorable effect on the plant. Do not eat, drink, or smoke during application; wash your hands before and after using the product.







FULVIC PROTECT FULVIC ACIDS AT 80 %



SOIL IMPROVER

APPLICATION FREQUENCY

It is recommended to make 2-3 applications per hectare for an optimal colonization of the plant with a frequency of 7-10 days each, which will achieve in the plant a healthy nutrition and defense of the crop.

DOSE

Crop	General
Dose	250g/ha
Frequency	Monthly
Application	2-3 applications

COMPATIBILITY WITH OTHER PRODUCTS

FulvicProtect is compatible with some fertilizers. In case of applying a bactericide, fungicide, pesticide or insecticide, it is necessary to check the residence time of said product and make sure that the application of **FulvicProtect** is outside this period, as well as wash the material with which the application is going to be made well with neutral soap to ensure that traces of the product are avoided. Consult your technical advisor for compatibility with other products. It is not recommended to apply **FulvicProtect** if a subsequent application of bactericides, fungicides or any similar product is planned as this would kill the population of microorganisms applied with **FulvicProtect**.

STORAGE AND TRANSPORT

Store the product in cool, covered places at room temperature. Do not expose it to the sun's rays.

ΤΟΧΙCΙΤΥ

100% organic product. It is not phytotoxic at the recommended concentrations, stages and forms of application. In case of ingestion or discomfort at the time of use, induce vomiting and consult your doctor.

DATA SHEET



If you think of microorganisms, think of Microendo.

www.microendo.com.mx @ info@microendo.com.mx @ 33 2603 8388 S

33 1469 5219 🔊